arXiv:1304.2577v1 [math.AP] 9 Apr 2013

On the Cauchy problem for a generalized Camassa-Holm equation with both quadratic and cubic nonlinearity

Xingxing Liu[1]* Zhijun Qiao[2]† and Zhaoyang Yin[1]‡
[1]Department of Mathematics, Sun Yat-sen University,
Guangzhou, Guangdong 510275 , China
[2]Department of Mathematics, University of Texas-Pan American,
Edinburg, Texas 78541, USA

Abstract

In this paper, we study the Cauchy problem for a generalized integrable Camassa-Holm equation with both quadratic and cubic nonlinearity, which has significant differences from the standard Camassa-Holm equation about the dynamics of the two-peakon and peakon-kink solutions (see [26] for details). We firstly establish the local well-posedness result in Besov spaces, and then present a precise blow-up scenario for strong solutions. Furthermore, we discuss weak kink solutions and their interactions with peakons.

2000 Mathematics Subject Classification: 35G25, 35L05

Keywords: Generalized Camassa-Holm equation, Besov spaces, local well-posedness, blow-up scenario.

1 Introduction

Recently, the following modified Camassa-Holm equation with cubic nonlinearity

$$m_t + [(u^2 - u_x^2)m]_x = 0, \quad m = u - u_{xx}, \tag{1.1}$$

has attracted attention in the literature [2, 13]. Eq. (1.1) was derived independently by Fokas (1995) [10], by Fuchssteiner (1996) [12], by Olver and Rosenau (1996) [22], and by Qiao (2006) [24]. Thanks to the work done in [24] about its generation from the two-dimensional Euler equation, Lax pair, M/W-shaped soliton, and peaked/cusped solitons, Eq. (1.1) regains remarkable

---

*E-mail: liuxingxing123456@163.com
†E-mail: qiao@utpa.edu
‡E-mail: mcsyzy@mail.sysu.edu.cn

attention in the study of peakon and integrable cubic nonlinear system ten years after they [10, 12, 22] presented it as a generalization of integrable system by applying the general method of tri-Hamiltonian duality to the bi-Hamiltonian representation of the classically integrable system. In fact, the basic method "Hamiltonian duality" used in [10, 11, 12, 22] has already given rise to some new bi-Hamiltonian equations. For example, the classical Camassa-Holm (CH) equation [3, 4]

$$m_t + um_x + 2mu_x = 0, \quad m = u - u_{xx},$$

can be derived from the Korteweg-de Vries (KdV) equation by tri-Hamiltonian duality, which describes the unidirectional propagation of waves at the free surface of shallow water under the influence of gravity. The variables $u(t,x)$ and $m(t,x)$ represent the velocity and its potential density of the fluid, respectively. In the past few years, a lot of literature has devoted to the investigation of the CH equation, because it has both peaked soliton (peakon) solutions and wave breaking phenomena, *i.e.*, the solution remains bounded while the slope of $u(t,x)$ becomes unbounded in finite time [5, 6, 7, 8, 17, 19]. However, by applying the "Hamiltonian duality" method to the modified Korteweg-de Vries (mKdV) equation, it is exactly Eq. (1.1). In contrast to the CH equation, Eq. (1.1) has nonlinear terms that are cubic, rather than quadratic. Another integrable CH type equation with cubic nonlinearity is the so-called Novikov equation [21]. The bi-Hamiltonian structure, complete integrability, the existence of peakon solutions, local well-posedness, the existence of low regularity solutions to the Novikov equation have been investigated extensively in [14, 15, 16, 20, 27, 28, 29].

Eq. (1.1) was recognized as an integrable nonlinear water wave model since it can be derived from the two-dimensional Euler equation and has Lax pair [24], which may allow it to be solved by the inverse scattering transform (IST) method [24, 25]. Moreover, Eq. (1.1) was found to admit the M/W-shaped soliton solutions [18, 24], which were not presented in the related study of CH equation [17]. In [13], Gui *et al.* considered the formation of singularities to Eq. (1.1), and singularities of the solutions could occur only in the form of wave-breaking. Then they discussed the existence of single and multi-peakon solutions for Eq. (1.1). In [2], Bies *et al.* called Eq. (1.1) the modified Korteweg-de Vries-Fokas-Qiao equation and used the geometric and analytic approaches to study it, and gave a geometric interpretation to the variable $m(t,x)$ and the Lie-algebraic structure of symmetries of Eq. (1.1). Then they applied the method in [19] to show the existence and uniqueness of weak solutions in the Sobolev space $H^s(\mathbb{R}), s > \frac{5}{2}$.

In the present paper, we consider the Cauchy problem of partial differential equation with both quadratic and cubic nonlinearity, which was recently proposed by Qiao, Xia and Li [26]

$$\begin{cases} m_t = \frac{1}{2}k_1[(u^2 - u_x^2)m]_x + \frac{1}{2}k_2(um_x + 2mu_x), & t > 0, x \in \mathbb{R}, \\ u(0,x) = u_0(x), \quad m = u - u_{xx}, & x \in \mathbb{R}, \end{cases} \tag{1.2}$$

where we usually take k_1, k_2 as non-positive constants. Indeed, Eq. (1.2) is a linear combination of CH equation and Eq. (1.1). Obviously, Eq. (1.2) is exactly the CH equation for $k_1 = 0, k_2 = -2$. When $k_1 = -2, k_2 = 0$, Eq. (1.2) reduces to the cubic nonlinear Eq. (1.1). Inspired by Gardner equation (known as a linear combination of KdV equation and mKdV equation [9, 23]), and derivation of the CH equation and Eq. (1.1) from the KdV equation and the mKdV equation by the "Hamiltonian duality" method, Qiao, Xia, and Li [26] study Eq. (1.2) and present the Lax pair and bi-Hamiltonian structure to Eq. (1.2), *i.e*,

$$m_t = J\frac{\delta H_1}{\delta m} = K\frac{\delta H_2}{\delta m},$$

where

$$J = k_1\partial m\partial^{-1}m\partial + \frac{1}{2}k_2(\partial m + m\partial), \quad K = \partial - \partial^3,$$

and two Hamiltonians are

$$H_1 = \frac{1}{2}\int_{\mathbb{R}}(u^2 + u_x^2)dx,$$

and

$$H_2 = \frac{1}{8}\int_{\mathbb{R}}(k_1u^4 + 2k_1u^2u_x^2 - \frac{1}{3}k_1u_x^4 + 2k_2u^3 + 2k_2uu_x^2)dx.$$

Thus, Eq. (1.2) is completely integrable.

Motivated by the study of the CH equation [8], our main work is to prove the local well-posedness to the Cauchy problem (1.2) in the nonhomogeneous Besov spaces. However, one of the differences is that we should deal with cubic nonlinearity in Besov spaces. Moreover, the nonlinear term "$m_xu_x^2$" makes us have to solve a transport equation satisfied by m, rather than u. In contrast to the case of the CH equation with initial data u_0 in the Sobolev space $H^s(\mathbb{R}), s > \frac{3}{2}$, we can only prove the well-posedness result with the initial profile u_0 in $H^s(\mathbb{R}), s > \frac{5}{2}$. In our procedure, we have overcome the critical index case by the interpolation method when we apply the transport theory to Eq. (1.2). Another one of the differences is that Eq. (1.2) possesses both quadratic and cubic nonlinearity. To get the uniform boundedness of the approximate solutions $\{u^{(n)}\}_{n\in\mathbb{N}}$, we have to handle the quadratic and cubic nonlinear terms together in Eq. (1.2). To overcome these difficulties, we consider two cases: the small initial data and the large one. With the local well-posedness result, we may naturally present a precise blow-up scenario to Eq. (1.2) by combining the blow-up criterion of the CH equation and the one of Eq. (1.1).

The entire paper is organized as follows. In Section 2, we present some facts on Besov spaces, some preliminary properties and the transport equation theory. In Section 3, we establish the local well-posedness result of Eq. (1.2) in Besov spaces. In Section 4, we derive a blow-up scenario for strong solutions to Eq. (1.2). In Section 5, we provide peakon and weak kink solutions and their interactions in an explicit form.

Notation. In the following, we denote $C > 0$ a generic constant only depending on p, r, s. Since our discussion about Eq. (1.2) is mainly on the line $\mathbb{R}$, for simplicity, we omit $\mathbb{R}$ in our notations of function spaces. And we denote the Fourier transform of a function u as $\mathcal{F}u$.

2 Preliminaries

In this section, we will recall some basic theory of the Littlewood-Paley decomposition and the transport equation theory on Besov spaces, which will play an important role in the sequel. One may get more details from [1, 8].

Proposition 2.1. *[1] (Littlewood-Paley decomposition) Let* $\mathcal{B} := \{\xi \in \mathbb{R}, |\xi| \leq \frac{4}{3}\}$ *and* $\mathcal{C} := \{\xi \in \mathbb{R}, \frac{3}{4} \leq |\xi| \leq \frac{8}{3}\}$*. Then there exist* $\psi(\xi) \in C_c^\infty(\mathcal{B})$ *and* $\varphi(\xi) \in C_c^\infty(\mathcal{C})$ *such that*

$$\psi(\xi) + \sum_{q\in\mathbb{N}} \varphi(2^{-q}\xi) = 1, \quad \forall\ \xi \in \mathbb{R}.$$

and

$$Supp\varphi(2^{-q}\cdot) \cap Supp\varphi(2^{-q'}\cdot) = \emptyset, \quad if\ |q-q'| \geq 2,$$
$$Supp\psi(\cdot) \cap Supp\varphi(2^{-q}\cdot) = \emptyset, \quad if\ q \geq 1.$$

Then for all $u \in \mathcal{S}'$ *(*$\mathcal{S}'$ *denotes the tempered distribution spaces), we can define the nonhomogeneous Littlewood-Paley decomposition of a distribution* u*.*

$$u = \sum_{q\in\mathbb{Z}} \Delta_q u,$$

where the localization operators are defined as follows:

$$\Delta_q u := 0, \quad for\ q \leq -2, \quad \Delta_{-1} u := \psi(D)u = \mathcal{F}^{-1}(\psi\mathcal{F}u),$$

and

$$\Delta_q u := \varphi(2^{-q}D)u = \mathcal{F}^{-1}(\varphi(2^{-q}\xi)\mathcal{F}u), \quad for\ q \geq 0.$$

Furthermore, we can define the low frequency cut-off operator S_q *as follows:*

$$S_q u := \sum_{i=-1}^{q-1} \Delta_i u = \psi(2^{-q}D)u = \mathcal{F}^{-1}(\psi(2^{-q}\xi)\mathcal{F}u).$$

Definition 2.1. *[1] (Besov spaces) Let* $s \in \mathbb{R}$*,* $1 \leq p, r \leq \infty$*. The nonhomogeneous Besov space* $B^s_{p,r}(\mathbb{R})$ *(*$B^s_{p,r}$ *for short) is defined by*

$$B^s_{p,r} := \{u \in \mathcal{S}'(\mathbb{R});\ \|u\|_{B^s_{p,r}} < \infty\},$$

where

$$\|u\|_{B^s_{p,r}} := \begin{cases} (\sum\limits_{q\in\mathbb{Z}} 2^{qsr}\|\Delta_q u\|^r_{L^p})^{\frac{1}{r}}, & r < \infty, \\ \sup\limits_{q\in\mathbb{Z}} 2^{qs}\|\Delta_q u\|_{L^p}, & r = \infty. \end{cases}$$

If $s = \infty$*,* $B^\infty_{p,r} := \bigcap\limits_{s\in\mathbb{R}} B^s_{p,r}$*.*

In order to state the local well-posedness result, we need to define the following spaces.

Definition 2.2. *Let $T>0, s\in\mathbb{R}$ and $1\le p\le\infty$. We define*

$$E^s_{p,r}(T) := C([0,T];B^s_{p,r})\cap C^1([0,T];B^{s-1}_{p,r}), \quad \text{for } r<\infty,$$

$$E^s_{p,\infty}(T) := L^\infty([0,T];B^s_{p,\infty})\cap Lip([0,T];B^{s-1}_{p,\infty}),$$

and

$$E^s_{p,r} := \bigcap_{T>0} E^s_{p,r}(T).$$

Next, we list the following useful properties for Besov spaces.

Proposition 2.2. *[1, 8] Let $s\in\mathbb{R}$, $1\le p,r,p_i,r_i\le\infty, i=1,2$. Then*
(i) Density: if $1\le p,r<\infty$, then $\mathcal{C}^\infty_c$ is dense in $B^s_{p,r}$.
(ii) Embedding: $B^s_{p_1,r_1}\hookrightarrow B^{s-(\frac{1}{p_1}-\frac{1}{p_2})}_{p_2,r_2}$, for $p_1\le p_2$ and $r_1\le r_2$,

$$B^{s_2}_{p,r_2}\hookrightarrow B^{s_1}_{p,r_1} \text{ locally compact}, \quad \text{for } s_1<s_2.$$

(iii) Algebraic properties: if $s>0$, $B^s_{p,r}\cap L^\infty$ is an algebra. Furthermore, $B^s_{p,r}$ is an algebra, provided that $s>\frac{1}{p}$ or $s\ge\frac{1}{p}$ and $r=1$.
(iv) Fatou lemma: if $\{u^{(n)}\}_{n\in\mathbb{N}}$ is bounded in $B^s_{p,r}$ and tends to u in $\mathcal{S}'$, then $u\in B^s_{p,r}$. Moreover,

$$\|u\|_{B^s_{p,r}}\le\liminf_{n\to\infty}\|u^{(n)}\|_{B^s_{p,r}}.$$

(v) Complex interpolation: if $u\in B^{s_1}_{p,r}\cap B^{s_2}_{p,r}$, then for all $\theta\in[0,1]$, we have $u\in B^{\theta s_1+(1-\theta)s_2}_{p,r}$. Moreover,

$$\|u\|_{B^{\theta s_1+(1-\theta)s_2}_{p,r}}\le\|u\|^\theta_{B^{s_1}_{p,r}}\|u\|^{(1-\theta)}_{B^{s_2}_{p,r}}.$$

(vi) One-dimensional Morse-type estimate:
1) If $s>0$,

$$\|uv\|_{B^s_{p,r}}\le C(\|u\|_{B^s_{p,r}}\|v\|_{L^\infty}+\|u\|_{L^\infty}\|v\|_{B^s_{p,r}}).$$

2) If $s_1\le\frac{1}{p}$, $s_2>\frac{1}{p}(s_2\ge\frac{1}{p}$ if $r=1)$ and $s_1+s_2>0$,

$$\|uv\|_{B^{s_1}_{p,r}}\le C\|u\|_{B^{s_1}_{p,r}}\|v\|_{B^{s_2}_{p,r}},$$

where C is a constant independent of u and v.
(vii) Action of Fourier multipliers on Besov spaces: let $m\in\mathbb{R}$ and f be a S^m-multiplier (i.e., $f:\mathbb{R}\to\mathbb{R}$ is a smooth function and satisfies that for each multi-index α, there exists a constant C_α such that $|\partial^\alpha f(\xi)|\le C_\alpha(1+|\xi|)^{m-|\alpha|}$, for $\forall\xi\in\mathbb{R}$.) Then the operator $f(D)$ is continuous from $B^s_{p,r}$ to $B^{s-m}_{p,r}$.

Now we state the following transport equation theory that is crucial to prove local well-posedness for Eq. (1.2).

Lemma 2.1. *[1, 8] (A priori estimate) Let* $1 \leq p, r \leq +\infty$ *and* $s > -\min\{\frac{1}{p}, 1-\frac{1}{p}\}$. *Assume that* v *be a function such that* $\partial_x v$ *belongs to* $L^1([0,T]; B^{s-1}_{p,r})$ *if* $s > 1+\frac{1}{p}$ *or to* $L^1([0,T]; B^{\frac{1}{p}}_{p,r} \cap L^\infty)$ *otherwise. Suppose also that* $f_0 \in B^s_{p,r}$, $F \in L^1([0,T]; B^s_{p,r})$, *and that* $f \in L^\infty([0,T]; B^s_{p,r}) \cap C([0,T]; \mathcal{S}')$ *be the solution of the one-dimensional transport equation*

$$\begin{cases} \partial_t f + v \cdot \partial_x f = F, \\ f \mid_{t=0} = f_0. \end{cases} \tag{2.1}$$

Then there exists a constant C *depending only on* s, p, r *such that the following statements hold for* $t \in [0,T]$
(i) If $r = 1$ *or* $s \neq 1 + \frac{1}{p}$,

$$\|f(t)\|_{B^s_{p,r}} \leq \|f_0\|_{B^s_{p,r}} + \int_0^t \|F(\tau)\|_{B^s_{p,r}} d\tau + C \int_0^t V'(\tau) \|f(\tau)\|_{B^s_{p,r}} d\tau,$$

or

$$\|f(t)\|_{B^s_{p,r}} \leq e^{CV(t)} (\|f_0\|_{B^s_{p,r}} + \int_0^t e^{-CV(\tau)} \|F(\tau)\|_{B^s_{p,r}} d\tau),$$

where

$$V(t) = \begin{cases} \int_0^t \|\partial_x v(\tau, \cdot)\|_{(B^{\frac{1}{p}}_{p,r} \cap L^\infty)} d\tau, & s < 1 + \frac{1}{p}, \\ \int_0^t \|\partial_x v(\tau, \cdot)\|_{B^{s-1}_{p,r}} d\tau, & s > 1 + \frac{1}{p}. \end{cases}$$

(ii) If $s \leq 1 + \frac{1}{p}$, *and* $\partial_x f_0, \partial_x f \in L^\infty([0,T] \times \mathbb{R})$ *and* $\partial_x F \in L^1([0,T]; L^\infty)$, *then*

$$\begin{aligned} & \|f(t)\|_{B^s_{p,r}} + \|\partial_x f(t)\|_{L^\infty} \\ \leq \; & e^{CV(t)} (\|f_0\|_{B^s_{p,r}} + \|\partial_x f_0\|_{L^\infty} + \int_0^t e^{-CV(\tau)} (\|F(\tau)\|_{B^s_{p,r}} + \|\partial_x F(\tau)\|_{L^\infty}) d\tau), \end{aligned}$$

where $V(t) = \int_0^t \|\partial_x v(\tau, \cdot)\|_{(B^{\frac{1}{p}}_{p,r} \cap L^\infty)} d\tau$.
(iii) If $f = v$, *then for all* $s > 0$, *the estimate in (i) holds with* $V(t) = \int_0^t \|\partial_x v(\tau, \cdot)\|_{L^\infty} d\tau$.
(iv) If $r < \infty$, *then* $f \in C([0,T]; B^s_{p,r})$. *If* $r = \infty$, *then* $f \in C([0,T]; B^{s'}_{p,1})$ *for all* $s' < s$.

Lemma 2.2. *[8] (Existence and uniqueness) Let* p, r, s, f_0 *and* F *be as in the statement of Lemma 2.1. Suppose that* $v \in L^\rho([0,T]; B^{-M}_{\infty,\infty})$ *for some* $\rho > 1, M > 0$ *and* $\partial_x v \in L^1([0,T]; B^{\frac{1}{p}}_{p,\infty} \cap L^\infty)$ *if* $s < 1 + \frac{1}{p}$, *and* $\partial_x v \in L^1([0,T]; B^{s-1}_{p,r})$ *if* $s > 1 + \frac{1}{p}$ *or* $s = 1 + \frac{1}{p}$ *and* $r = 1$. *Then the transport equation (2.1) has a unique solution* $f \in L^\infty([0,T]; B^s_{p,r}) \bigcap (\bigcap_{s'<s} C([0,T]; B^{s'}_{p,1}))$ *and the corresponding inequalities in Lemma 2.1 hold true. Moreover, if* $r < \infty$, *then* $f \in C([0,T]; B^s_{p,r})$.

3 Local well-posedness

In this section, we shall study the local well-posedness of Eq.(1.2) in the nonhomogeneous Besov spaces. At first, we present a priori estimates about the solutions of Eq.(1.2), which can be applied to prove the uniqueness and continuity with the initial data in some sense.

Lemma 3.1. *Suppose that* $1 \leq p, r \leq \infty$ *and* $s > \max\{2+\frac{1}{p}, \frac{5}{2}, 3-\frac{1}{p}\}$. *Let* $u^{(1)}, u^{(2)} \in L^\infty([0,T]; B^s_{p,r}) \cap C([0,T]; \mathcal{S}')$ *be two given solutions to Eq.(1.2) with initial data* $u^{(1)}_0, u^{(2)}_0 \in B^s_{p,r}$, *and let* $u^{(12)} := u^{(2)} - u^{(1)}$ *and* $m^{(12)} := m^{(2)} - m^{(1)}$. *Then for all* $t \in [0,T]$, *we have*
(1) If $s > \max\{2+\frac{1}{p}, \frac{5}{2}, 3-\frac{1}{p}\}$ *and* $s \neq 4+\frac{1}{p}$, *then*

$$\begin{aligned} \|u^{(12)}\|_{B^{s-1}_{p,r}} \leq\ & \|u^{(12)}_0\|_{B^{s-1}_{p,r}} \exp\{C\int_0^t (\|u^{(1)}(\tau)\|^2_{B^s_{p,r}} + \|u^{(2)}(\tau)\|^2_{B^s_{p,r}} \\ & + \|u^{(1)}(\tau)\|_{B^s_{p,r}} + \|u^{(2)}(\tau)\|_{B^s_{p,r}}) d\tau\}. \end{aligned}$$

(2) If $s = 4+\frac{1}{p}$, *then*

$$\begin{aligned} & \|u^{(12)}\|_{B^{s-1}_{p,r}} \\ \leq\ & C\|u^{(12)}_0\|^\theta_{B^{s-1}_{p,r}} (\|u^{(1)}\|_{B^s_{p,r}} + \|u^{(2)}\|_{B^s_{p,r}})^{1-\theta} \exp\{\theta C\int_0^t (\|u^{(1)}(\tau)\|^2_{B^s_{p,r}} \\ & + \|u^{(2)}(\tau)\|^2_{B^s_{p,r}} + \|u^{(1)}(\tau)\|_{B^s_{p,r}} + \|u^{(2)}(\tau)\|_{B^s_{p,r}}) d\tau\}, \end{aligned}$$

with $\theta = \frac{1}{2}(1-\frac{1}{2p}) \in (0,1)$.

Proof. It is obvious that $u^{(12)} \in L^\infty([0,T]; B^s_{p,r}) \cap C([0,T]; \mathcal{S}')$ and $u^{(12)}, m^{(12)}$ solves the following transport equation

$$\begin{cases} m^{(12)}_t - \{\frac{k_1}{2}[(u^{(1)})^2 - (u^{(1)}_x)^2] + \frac{k_2}{2}u^{(1)}\} m^{(12)}_x = F(t,x), & t > 0, x \in \mathbb{R}, \\ m^{(12)}\mid_{t=0} = m^{(12)}_0 := m^{(2)}_0 - m^{(1)}_0, \quad m = u - u_{xx}, & x \in \mathbb{R}, \end{cases} \tag{3.1}$$

where $F(t,x) := \frac{k_1}{2}[u^{(12)}(u^{(1)} + u^{(2)}) - u^{(12)}_x(u^{(1)}_x + u^{(2)}_x)]\partial_x m^{(2)} + k_1[u^{(12)}_x (m^{(2)})^2$ $+ u^{(1)}_x m^{(12)}(m^{(1)} + m^{(2)})] + \frac{k_2}{2}u^{(12)} m^{(2)}_x + k_2 u^{(12)}_x m^{(2)} + k_2 u^{(1)}_x m^{(12)}$.

Applying Lemma 2.1 to the transport equation (3.1), we have

$$\begin{aligned} \|m^{(12)}\|_{B^{s-3}_{p,r}} \leq\ & \|m^{(12)}_0\|_{B^{s-3}_{p,r}} + \int_0^t \|F(\tau)\|_{B^{s-3}_{p,r}} d\tau \\ & + C\int_0^t \|(u^{(1)})^2 - (u^{(1)}_x)^2 + u^{(1)}\|_{B^{s-3}_{p,r}} \|m^{(12)}\|_{B^{s-3}_{p,r}} d\tau. \end{aligned} \tag{3.2}$$

Indeed, if $\max\{2+\frac{1}{p}, \frac{5}{2}\} < s \leq 3+\frac{1}{p}$, by Proposition 2.2 (vi), we get

$$\begin{aligned}
& \|F(\tau)\|_{B^{s-3}_{p,r}} \\
\leq\ & C\{\|[u^{(12)}(u^{(1)}+u^{(2)}) - u_x^{(12)}(u_x^{(1)}+u_x^{(2)})]\partial_x m^{(2)}\|_{B^{s-3}_{p,r}} \\
& +\|u_x^{(12)}(m^{(2)})^2\|_{B^{s-3}_{p,r}} + \|u_x^{(1)}m^{(12)}(m^{(1)}+m^{(2)})\|_{B^{s-3}_{p,r}} \\
& +\|u^{(12)}m_x^{(2)} + u_x^{(12)}m^{(2)} + u_x^{(1)}m^{(12)}\|_{B^{s-3}_{p,r}}\} \\
\leq\ & C\{\|m_x^{(2)}\|_{B^{s-3}_{p,r}}\|[u^{(12)}(u^{(1)}+u^{(2)}) - u_x^{(12)}(u_x^{(1)}+u_x^{(2)})]\|_{B^{s-2}_{p,r}} \\
& +\|u_x^{(12)}\|_{B^{s-3}_{p,r}}\|(m^{(2)})^2\|_{B^{s-2}_{p,r}} + \|m^{(12)}\|_{B^{s-3}_{p,r}}\|u_x^{(1)}(m^{(1)}+m^{(2)})\|_{B^{s-2}_{p,r}} \\
& +\|m_x^{(2)}\|_{B^{s-3}_{p,r}}\|u^{(12)}\|_{B^{s-2}_{p,r}} + \|u_x^{(12)}\|_{B^{s-3}_{p,r}}\|m^{(2)}\|_{B^{s-2}_{p,r}} \\
& +\|m^{(12)}\|_{B^{s-3}_{p,r}}\|u_x^{(1)}\|_{B^{s-2}_{p,r}}\}.
\end{aligned}$$

Since $s > \max\{2+\frac{1}{p}, \frac{5}{2}\}$, we know that $B^{s-2}_{p,r}$ is an algebra. Thus, we deduce

$$\begin{aligned}
& \|F(\tau)\|_{B^{s-3}_{p,r}} \\
\leq\ & C[\|u^{(12)}\|_{B^{s-1}_{p,r}}(\|u^{(1)}\|^2_{B^s_{p,r}} + \|u^{(2)}\|^2_{B^s_{p,r}} + \|u^{(1)}\|_{B^s_{p,r}} + \|u^{(2)}\|_{B^s_{p,r}}).
\end{aligned} \tag{3.3}$$

For $s > 3+\frac{1}{p}$, the inequality (3.3) also holds true in view of the fact that $B^{s-3}_{p,r}$ is an algebra. Note that

$$\begin{aligned}
& \|(u^{(1)})^2 - (u_x^{(1)})^2 + u^{(1)}\|_{B^{s-3}_{p,r}}\|m^{(12)}\|_{B^{s-3}_{p,r}} \\
\leq\ & C[\|u^{(12)}\|_{B^{s-1}_{p,r}}(\|u^{(1)}\|^2_{B^s_{p,r}} + \|u^{(2)}\|^2_{B^s_{p,r}} + \|u^{(1)}\|_{B^s_{p,r}} + \|u^{(2)}\|_{B^s_{p,r}}).
\end{aligned}$$

Therefore, inserting the above inequality and (3.3) into (3.2), we obtain

$$\begin{aligned}
\|u^{(12)}\|_{B^{s-1}_{p,r}} \leq\ & \|u_0^{(12)}\|_{B^{s-1}_{p,r}} + C\int_0^t [\|u^{(12)}(\tau)\|_{B^{s-1}_{p,r}} \\
& \times(\|u^{(1)}\|^2_{B^s_{p,r}} + \|u^{(2)}\|^2_{B^s_{p,r}} + \|u^{(1)}\|_{B^s_{p,r}} + \|u^{(2)}\|_{B^s_{p,r}})]d\tau.
\end{aligned}$$

Then, by Gronwall's inequality, we prove (1).

Since we can not apply Lemma 2.1 to (3.1) for the critical case $s = 4+\frac{1}{p}$, we here use the interpolation method to deal with it.

In fact, we can choose $\theta = \frac{1}{2}(1-\frac{1}{2p}) \in (0,1)$, such that $s-1 = 3+\frac{1}{p} = (1-\theta)(4+\frac{1}{2p})+\theta(2+\frac{1}{2p})$. Then, by Proposition 2.2 (v), we have

$$\|u^{(12)}\|_{B^{s-1}_{p,r}} = \|u^{(12)}\|_{B^{3+\frac{1}{p}}_{p,r}} \leq \|u^{(12)}\|^{\theta}_{B^{2+\frac{1}{2p}}_{p,r}}\|u^{(12)}\|^{1-\theta}_{B^{4+\frac{1}{2p}}_{p,r}}.$$

Then, from the obtained result of (1) in this lemma, we get

$$\begin{aligned}
&\|u^{(12)}\|_{B^{s-1}_{p,r}}\\
\leq\ & [\|u_0^{(12)}\|_{B^{2+\frac{1}{2p}}_{p,r}} \exp(C\int_0^t (\|u^{(1)}\|^2_{B^{3+\frac{1}{2p}}_{p,r}} + \|u^{(2)}\|^2_{B^{3+\frac{1}{2p}}_{p,r}}\\
&+\|u^{(1)}\|_{B^{3+\frac{1}{2p}}_{p,r}} + \|u^{(2)}\|_{B^{3+\frac{1}{2p}}_{p,r}})d\tau)]^{\theta} \times (\|u^{(1)}\|_{B^{4+\frac{1}{2p}}_{p,r}} + \|u^{(2)}\|_{B^{4+\frac{1}{2p}}_{p,r}})^{1-\theta}\\
\leq\ & C\|u_0^{(12)}\|^{\theta}_{B^{s-1}_{p,r}}(\|u^{(1)}\|_{B^s_{p,r}} + \|u^{(2)}\|_{B^s_{p,r}})^{1-\theta} \exp\{\theta C\int_0^t (\|u^{(1)}(\tau)\|^2_{B^s_{p,r}}\\
&+\|u^{(2)}(\tau)\|^2_{B^s_{p,r}} + \|u^{(1)}(\tau)\|_{B^s_{p,r}} + \|u^{(2)}(\tau)\|_{B^s_{p,r}})d\tau\}.
\end{aligned}$$

This completes the proof of Lemma 3.1. □

Next, we use the classical Friedrichs regularization method to construct the approximation solutions to Eq. (1.2).

Lemma 3.2. *Suppose that p, r and s be as in the statement of Lemma 3.1, $u_0 \in B^s_{p,r}$ and $u^{(0)} := 0$. Then there exists a sequence of smooth functions $\{u^{(n)}\}_{n\in\mathbb{N}} \in C(\mathbb{R}^+; B^{\infty}_{p,r})$ solving the following transport equation by induction*

$$\begin{cases}
m_t^{(n+1)} - \{\frac{k_1}{2}[(u^{(n)})^2 - (u_x^{(n)})^2] + \frac{k_2}{2}u^{(n)}\}m_x^{(n+1)} = \\
\qquad\qquad k_1 u_x^{(n)}(m^{(n)})^2 + k_2 u_x^{(n)} m^{(n)}, & t > 0, x \in \mathbb{R},\\
u^{(n+1)}\mid_{t=0} = u_0^{(n+1)}(x) = S_{n+1}u_0, & x \in \mathbb{R}.
\end{cases} \tag{3.4}$$

Moreover, there exists a $T > 0$ such that the solutions satisfying the following properties.
(1) $\{u^{(n)}\}_{n\in\mathbb{N}}$ is uniformly bounded in $E^s_{p,r}(T)$.
(2) $\{u^{(n)}\}_{n\in\mathbb{N}}$ is a Cauchy sequence in $C([0,T]; B^{s-1}_{p,r})$.

Proof. By the definition of S_q, we know that all the data $S_{n+1}u_0 \in B^{\infty}_{p,r}$. Thus, from Lemma 2.2, we deduce by induction that for all $n \in \mathbb{N}$, Eq. (3.4) has a global solution belonging $C(\mathbb{R}^+; B^{\infty}_{p,r})$.

For $s > \max\{2 + \frac{1}{p}, \frac{5}{2}, 3 - \frac{1}{p}\}$ and $s \neq 4 + \frac{1}{p}$, by Lemma 2.1, we obtain

$$\begin{aligned}
\|m^{(n+1)}\|_{B^{s-2}_{p,r}} \leq\ & e^{C\int_0^t \|[(u^{(n)})^2 - (u_x^{(n)})^2 + u^{(n)}](\tau)\|_{B^{s-2}_{p,r}} d\tau}(\|m_0\|_{B^{s-2}_{p,r}}\\
&+C\int_0^t e^{-C\int_0^{\tau} \|[(u^{(n)})^2 - (u_x^{(n)})^2 + u^{(n)}](\tau')\|_{B^{s-2}_{p,r}} d\tau'}\\
&\times\|u_x^{(n)}(m^{(n)})^2 + u_x^{(n)} m^{(n)}\|_{B^{s-2}_{p,r}} d\tau).
\end{aligned} \tag{3.5}$$

Since $s > 2 + \frac{1}{p}$, we know that $B^{s-2}_{p,r}$ is an algebra. Thus, we have

$$\begin{aligned}
&\|u_x^{(n)}(m^{(n)})^2 + u_x^{(n)} m^{(n)}\|_{B^{s-2}_{p,r}}\\
\leq\ & C\|u_x^{(n)}\|_{B^{s-2}_{p,r}}(\|(m^{(n)})^2\|_{B^{s-2}_{p,r}} + \|(m^{(n)})\|_{B^{s-2}_{p,r}})\\
\leq\ & C(\|u^{(n)}\|^3_{B^s_{p,r}} + \|u^{(n)}\|^2_{B^s_{p,r}}),
\end{aligned}$$

and

$$\|(u^{(n)})^2-(u_x^{(n)})^2+u^{(n)}\|_{B^{s-2}_{p,r}}\leq C(\|u^{(n)}\|^2_{B^s_{p,r}}+\|u^{(n)}\|_{B^s_{p,r}}).$$

Inserting the above inequalities into (3.5), we obtain

$$\begin{aligned}\|u^{(n+1)}\|_{B^s_{p,r}} &\leq e^{C\int_0^t(\|u^{(n)}\|^2_{B^s_{p,r}}+\|u^{(n)}\|_{B^s_{p,r}})(\tau)d\tau}\|u_0\|_{B^s_{p,r}}\\ &\quad +C\int_0^t e^{C\int_\tau^t(\|u^{(n)}\|^2_{B^s_{p,r}}+\|u^{(n)}\|_{B^s_{p,r}})(\tau')d\tau'}\\ &\quad \times(\|u^{(n)}\|^3_{B^s_{p,r}}+\|u^{(n)}\|^2_{B^s_{p,r}})d\tau. \qquad (3.6)\end{aligned}$$

In order to prove the uniform boundedness of $\{u^{(n)}\}_{n\in\mathbb{N}}$, we shall divide our discussion into two parts. When $2\|u_0\|_{B^s_{p,r}}<1$, we choose a $T_1>0$ such that

$$T_1<\min\{\frac{1-2\|u_0\|_{B^s_{p,r}}}{8C\|u_0\|_{B^s_{p,r}}},\frac{1}{4C}\},$$

and suppose by induction that for all $t\in[0,T_1]$

$$\|u^{(n)}\|_{B^s_{p,r}}\leq\frac{2\|u_0\|_{B^s_{p,r}}}{1-8C\|u_0\|_{B^s_{p,r}}t}. \qquad (3.7)$$

Noting that $t\leq T_1<\frac{1-2\|u_0\|_{B^s_{p,r}}}{8C\|u_0\|_{B^s_{p,r}}}$, we have $\frac{2\|u_0\|_{B^s_{p,r}}}{1-8C\|u_0\|_{B^s_{p,r}}t}<1$. By (3.7), we obtain

$$\begin{aligned}\|u^{(n)}\|_{B^s_{p,r}}&\leq\frac{2\|u_0\|_{B^s_{p,r}}}{1-8C\|u_0\|_{B^s_{p,r}}t}\\ &\leq(\frac{2\|u_0\|_{B^s_{p,r}}}{1-8C\|u_0\|_{B^s_{p,r}}t})^{\frac{2}{3}}\leq(\frac{2\|u_0\|_{B^s_{p,r}}}{1-8C\|u_0\|_{B^s_{p,r}}t})^{\frac{1}{2}}. \qquad (3.8)\end{aligned}$$

From (3.8), we can deduce that

$$\begin{aligned}&C\int_\tau^t(\|u^{(n)}\|^2_{B^s_{p,r}}+\|u^{(n)}\|_{B^s_{p,r}})(\tau')d\tau'\\ \leq\ &C\int_\tau^t\{[(\frac{2\|u_0\|_{B^s_{p,r}}}{1-8C\|u_0\|_{B^s_{p,r}}\tau'})^{\frac{1}{2}}]^2+\frac{2\|u_0\|_{B^s_{p,r}}}{1-8C\|u_0\|_{B^s_{p,r}}\tau'}\}d\tau'\\ \leq\ &-\frac{1}{2}\int_\tau^t\frac{-8C\|u_0\|_{B^s_{p,r}}}{1-8C\|u_0\|_{B^s_{p,r}}\tau'}d\tau'\\ =\ &\ln\sqrt{1-8C\|u_0\|_{B^s_{p,r}}\tau}-\ln\sqrt{1-8C\|u_0\|_{B^s_{p,r}}t}.\end{aligned}$$

Inserting the above inequality and (3.8) into (3.6) yields

$$\begin{aligned}
&\|u^{(n+1)}\|_{B^s_{p,r}} \\
\leq\ & \frac{\|u_0\|_{B^s_{p,r}}}{\sqrt{1-8C\|u_0\|_{B^s_{p,r}}t}}+\frac{C}{\sqrt{1-8C\|u_0\|_{B^s_{p,r}}t}}\{\int_0^t\sqrt{1-8C\|u_0\|_{B^s_{p,r}}\tau} \\
&\times[((\frac{2\|u_0\|_{B^s_{p,r}}}{1-8C\|u_0\|_{B^s_{p,r}}\tau})^{\frac{2}{3}})^3+((\frac{2\|u_0\|_{B^s_{p,r}}}{1-8C\|u_0\|_{B^s_{p,r}}\tau})^{\frac{1}{2}})^2]\}d\tau \\
\leq\ & \frac{\|u_0\|_{B^s_{p,r}}}{\sqrt{1-8C\|u_0\|_{B^s_{p,r}}t}}+\frac{\|u_0\|_{B^s_{p,r}}}{\sqrt{1-8C\|u_0\|_{B^s_{p,r}}t}}\int_0^t\frac{4C\|u_0\|_{B^s_{p,r}}}{(1-8C\|u_0\|_{B^s_{p,r}}\tau)^{\frac{3}{2}}}d\tau \\
&+\frac{1}{\sqrt{1-8C\|u_0\|_{B^s_{p,r}}t}}\int_0^t\frac{2C\|u_0\|_{B^s_{p,r}}}{(1-8C\|u_0\|_{B^s_{p,r}}\tau)^{\frac{1}{2}}}d\tau \\
\leq\ & \frac{\|u_0\|_{B^s_{p,r}}}{\sqrt{1-8C\|u_0\|_{B^s_{p,r}}t}}+\frac{\|u_0\|_{B^s_{p,r}}}{\sqrt{1-8C\|u_0\|_{B^s_{p,r}}t}}[(1-8C\|u_0\|_{B^s_{p,r}}\tau)^{-\frac{1}{2}}|_0^t] \\
&+\frac{1}{\sqrt{1-8C\|u_0\|_{B^s_{p,r}}t}}[-\frac{1}{2}(1-8C\|u_0\|_{B^s_{p,r}}\tau)^{\frac{1}{2}}|_0^t] \\
\leq\ & \frac{\|u_0\|_{B^s_{p,r}}}{\sqrt{1-8C\|u_0\|_{B^s_{p,r}}t}}(1+\frac{1}{\sqrt{1-8C\|u_0\|_{B^s_{p,r}}t}}-1)+\frac{1}{\sqrt{1-8C\|u_0\|_{B^s_{p,r}}t}} \\
&\times\frac{1}{2}(1-\sqrt{1-8C\|u_0\|_{B^s_{p,r}}t}) \\
\leq\ & \frac{2\|u_0\|_{B^s_{p,r}}}{1-8C\|u_0\|_{B^s_{p,r}}t},
\end{aligned}\tag{3.9}$$

where we used the following fact that

$$T_1<\frac{1}{4C}\Rightarrow\frac{1}{2}(1-\sqrt{1-8C\|u_0\|_{B^s_{p,r}}t})\leq\frac{\|u_0\|_{B^s_{p,r}}}{\sqrt{1-8C\|u_0\|_{B^s_{p,r}}t}},$$

in the last inequality. Thus, we prove (3.7) for the case $2\|u_0\|_{B^s_{p,r}}<1$.

On the other hand, when $2\|u_0\|_{B^s_{p,r}}\geq 1$, we choose a $T_2>0$ such that $T_2\leq\frac{1-e^{-1}}{16C\|u_0\|^2_{B^s_{p,r}}}<\frac{1}{16C\|u_0\|^2_{B^s_{p,r}}}$, and suppose by induction that for all $t\in[0,T_2]$

$$\|u^{(n)}\|_{B^s_{p,r}}\leq\frac{2\|u_0\|_{B^s_{p,r}}}{\sqrt{1-16C\|u_0\|^2_{B^s_{p,r}}t}}.\tag{3.10}$$

Noting that $2\|u_0\|_{B^s_{p,r}} \geq 1$, we get $\frac{2\|u_0\|_{B^s_{p,r}}}{(1-16C\|u_0\|^2_{B^s_{p,r}}t)^{\frac{1}{2}}} \geq 1$. From (3.10), we obtain

$$\begin{aligned}\|u^{(n)}\|_{B^s_{p,r}} &\leq \frac{2\|u_0\|_{B^s_{p,r}}}{\sqrt{1-16C\|u_0\|^2_{B^s_{p,r}}t}} \\ &\leq (\frac{2\|u_0\|_{B^s_{p,r}}}{\sqrt{1-16C\|u_0\|^2_{B^s_{p,r}}t}})^{\frac{3}{2}} \leq (\frac{2\|u_0\|_{B^s_{p,r}}}{\sqrt{1-16C\|u_0\|^2_{B^s_{p,r}}t}})^2. \end{aligned} \tag{3.11}$$

By (3.11), we find that

$$\begin{aligned} & C\int_\tau^t (\|u^{(n)}\|^2_{B^s_{p,r}} + \|u^{(n)}\|_{B^s_{p,r}})(\tau')d\tau' \\ \leq\ & C\int_\tau^t \{(\frac{2\|u_0\|_{B^s_{p,r}}}{\sqrt{1-16C\|u_0\|^2_{B^s_{p,r}}\tau'}})^2 + \frac{4\|u_0\|^2_{B^s_{p,r}}}{1-16C\|u_0\|^2_{B^s_{p,r}}\tau'}\}d\tau' \\ \leq\ & -\frac{1}{2}\int_\tau^t \frac{-16C\|u_0\|^2_{B^s_{p,r}}}{1-16C\|u_0\|^2_{B^s_{p,r}}\tau'}d\tau' \\ =\ & \ln\sqrt{1-16C\|u_0\|^2_{B^s_{p,r}}\tau} - \ln\sqrt{1-16C\|u_0\|^2_{B^s_{p,r}}t}. \end{aligned}$$

Inserting the above inequality and (3.11) into (3.6), we obtain

$$\begin{aligned} & \|u^{(n+1)}\|_{B^s_{p,r}} \\ \leq\ & \frac{\|u_0\|_{B^s_{p,r}}}{\sqrt{1-16C\|u_0\|^2_{B^s_{p,r}}t}} + \frac{C}{\sqrt{1-16C\|u_0\|^2_{B^s_{p,r}}t}}\{\int_0^t \sqrt{1-16C\|u_0\|^2_{B^s_{p,r}}\tau} \\ & \times[(\frac{2\|u_0\|_{B^s_{p,r}}}{\sqrt{1-16C\|u_0\|^2_{B^s_{p,r}}\tau}})^3 + ((\frac{2\|u_0\|_{B^s_{p,r}}}{\sqrt{1-16C\|u_0\|^2_{B^s_{p,r}}\tau}})^{\frac{3}{2}})^2]\}d\tau \\ \leq\ & \frac{\|u_0\|_{B^s_{p,r}}}{\sqrt{1-16C\|u_0\|^2_{B^s_{p,r}}t}} + \frac{\|u_0\|_{B^s_{p,r}}}{\sqrt{1-16C\|u_0\|^2_{B^s_{p,r}}t}}\int_0^t \frac{16C\|u_0\|^2_{B^s_{p,r}}}{(1-16C\|u_0\|^2_{B^s_{p,r}}\tau)}d\tau \\ \leq\ & \frac{\|u_0\|_{B^s_{p,r}}}{\sqrt{1-16C\|u_0\|^2_{B^s_{p,r}}t}}[1-\ln(1-16C\|u_0\|^2_{B^s_{p,r}}\tau)|_0^t] \\ \leq\ & \frac{2\|u_0\|_{B^s_{p,r}}}{\sqrt{1-16C\|u_0\|^2_{B^s_{p,r}}t}}, \end{aligned} \tag{3.12}$$

where we used the following fact that

$$T_2 \leq \frac{1-e^{-1}}{16C\|u_0\|^2_{B^s_{p,r}}} \Rightarrow 1-\ln(1-16C\|u_0\|^2_{B^s_{p,r}}t) \leq 2,$$

in the last inequality. Thus, we prove (3.10) for the case $2\|u_0\|_{B^s_{p,r}} \geq 1$.

Therefore, from the above discussion of the two cases, choosing $T = \min\{T_1, T_2\} > 0$, combining (3.9) and (3.12), we have proved that $\{u^{(n)}\}_{n\in\mathbb{N}}$ is uniformly bounded in $C([0,T]; B^s_{p,r})$. Using Proposition 2.2 (vi) and the fact $B^{s-2}_{p,r}$ is an algebra as $s > 2 + \frac{1}{p}$, we have

$$\begin{aligned} & \|[((u^{(n)})^2 - (u_x^{(n)})^2) + u^{(n)}]m_x^{(n+1)} + u_x^{(n)}(m^{(n)})^2 + u_x^{(n)}m^{(n)}\|_{B^{s-3}_{p,r}} \\ \leq\ & C\{\|m_x^{(n+1)}\|_{B^{s-3}_{p,r}}[(\|(u^{(n)})^2\|_{B^{s-2}_{p,r}} + \|(u_x^{(n)})^2\|_{B^{s-2}_{p,r}}) + \|u^{(n)}\|_{B^{s-2}_{p,r}}] \\ & +\|u_x^{(n)}\|_{B^{s-3}_{p,r}}(\|(m^{(n)})^2\|_{B^{s-2}_{p,r}} + \|m^{(n)}\|_{B^{s-2}_{p,r}})\} \\ \leq\ & C[\|u^{(n+1)}\|_{B^s_{p,r}}(\|(u^{(n)})\|^2_{B^s_{p,r}} + \|u^{(n)}\|_{B^s_{p,r}}) + (\|u^{(n)}\|^3_{B^s_{p,r}} + \|u^{(n)}\|^2_{B^s_{p,r}})]. \end{aligned}$$

From Eq. (3.4), we get $\partial_t m^{(n+1)} \in C \in ([0,T]; B^{s-3}_{p,r})$. Hence, $\partial_t u^{(n+1)} \in C \in ([0,T]; B^{s-1}_{p,r})$ is uniformly bounded, which yields that the sequence $\{u^{(n)}\}_{n\in\mathbb{N}}$ is uniformly bounded in $E^s_{p,r}(T)$.

Now it suffices to prove that

$$\{u^{(n)}\}_{n\in\mathbb{N}} \text{ is a Cauchy sequence in } C([0,T]; B^{s-1}_{p,r}).$$

Indeed, from Eq. (3.4), for all $n, l \in \mathbb{N}$, we have

$$\{\partial_t - [\frac{k_1}{2}((u^{(n+l)})^2 - (u_x^{(n+l)})^2) + \frac{k_2}{2}u^{(n+l)}]\}(m^{(n+l+1)} - m^{(n+1)}) = G(t,x),$$

where $G(t,x) := \{\frac{k_1}{2}[(u^{(n+l)} - u^{(n)})(u^{(n+l)} + u^{(n)}) - (u_x^{(n+l)} - u_x^{(n)})(u_x^{(n+l)} + u_x^{(n)})]$ $+ \frac{k_2}{2}(u^{(n+l)} - u^{(n)})\}\partial_x m^{(n+1)} + k_1 u_x^{(n+l)}(m^{(n+l)} - m^{(n)})(m^{(n+l)} + m^{(n)}) + k_1(u_x^{(n+l)} - u_x^{(n)})(m^{(n)})^2$ $+ k_2 u_x^{(n+l)}(m^{(n+l)} - m^{(n)}) + k_2(u_x^{(n+l)} - u_x^{(n)})m^{(n)}$.

Applying Lemma 2.1 again, for all $t \in [0,T]$, we have

$$\begin{aligned} & \|m^{(n+l+1)} - m^{(n+1)}\|_{B^{s-3}_{p,r}} \qquad (3.13) \\ \leq\ & e^{C\int_0^t \|[(u^{(n+l)})^2 - (u_x^{(n+l)})^2 + u^{(n+l)}](\tau)\|_{B^{s-2}_{p,r}} d\tau}(\|m_0^{(n+l+1)} - m_0^{(n+1)}\|_{B^{s-3}_{p,r}} \\ & +C\int_0^t e^{-C\int_0^\tau \|[(u^{(n+l)})^2 - (u_x^{(n+l)})^2 + u^{(n+l)}](\tau')\|_{B^{s-2}_{p,r}} d\tau'} \times \|G(\tau)\|_{B^{s-3}_{p,r}} d\tau). \end{aligned}$$

Similar to the proof of the estimate of $\|F(\tau)\|_{B^{s-3}_{p,r}}$ in Lemma 3.1, for $s > \max\{2 + \frac{1}{p}, \frac{5}{2}, 3 - \frac{1}{p}\}$ and $s \neq 4 + \frac{1}{p}$, we also obtain

$$\begin{aligned} \|G(\tau)\|_{B^{s-3}_{p,r}} \leq\ & C\|u^{(n+l)} - u^{(n)}\|_{B^{s-1}_{p,r}}(\|u^{(n)}\|^2_{B^s_{p,r}} + \|u^{(n+1)}\|^2_{B^s_{p,r}} \\ & +\|u^{(n+l)}\|^2_{B^s_{p,r}} + \|u^{(n)}\|_{B^s_{p,r}} + \|u^{(n+1)}\|_{B^s_{p,r}} + \|u^{(n+l)}\|_{B^s_{p,r}}). \end{aligned}$$

Inserting the above inequality into (3.13), we have

$$\begin{aligned}
& \|u^{(n+l+1)} - u^{(n+1)}\|_{B^{s-1}_{p,r}} \\
\leq\ & e^{C\int_0^t \|[(u^{(n+l)})^2-(u_x^{(n+l)})^2+u^{(n+l)}](\tau)\|_{B^{s-2}_{p,r}}d\tau}(\|u_0^{(n+l+1)} - u_0^{(n+1)}\|_{B^{s-1}_{p,r}} \\
& +C\int_0^t e^{-C\int_0^\tau \|[(u^{(n+l)})^2-(u_x^{(n+l)})^2+u^{(n+l)}](\tau')\|_{B^{s-2}_{p,r}}d\tau'}\|u^{(n+l)} - u^{(n)}\|_{B^{s-1}_{p,r}} \\
& \times(\|u^{(n)}\|^2_{B^s_{p,r}} + \|u^{(n+1)}\|^2_{B^s_{p,r}} + \|u^{(n+l)}\|^2_{B^s_{p,r}} + \|u^{(n)}\|_{B^s_{p,r}} + \|u^{(n+1)}\|_{B^s_{p,r}} \\
& +\|u^{(n+l)}\|_{B^s_{p,r}}).
\end{aligned}$$

Note that $\{u^{(n)}\}_{n\in\mathbb{N}}$ is uniformly bounded in $E^s_{p,r}(T)$ and

$$\begin{aligned}
& \|u_0^{(n+l+1)} - u_0^{(n+1)}\|_{B^{s-1}_{p,r}} \\
=\ & \|S_{n+l+1}u_0 - S_{n+1}u_0\|_{B^{s-1}_{p,r}} = \|\sum_{q=n+1}^{n+l}\Delta_q u_0\|_{B^{s-1}_{p,r}} \\
\leq\ & (\sum_{k\geq -1} 2^{k(s-1)r}\|\Delta_k(\sum_{q=n+1}^{n+l}\Delta_q u_0)\|^r_{L^p})^{\frac{1}{r}} \leq C(\sum_{k=n}^{n+l+1} 2^{-kr}2^{krs}\|\Delta_k u_0\|^r_{L^p})^{\frac{1}{r}} \\
\leq\ & C2^{-n}\|u_0\|_{B^s_{p,r}}.
\end{aligned}$$

Hence, there exists a constant C_T independent of n, l such that for all $t \in [0, T]$

$$\|(u^{(n+l+1)} - u^{(n+1)})(t)\|_{B^{s-1}_{p,r}} \leq C_T(2^{-n} + \int_0^t \|(u^{(n+l)} - u^{(n)})(\tau)\|_{B^{s-1}_{p,r}}d\tau).$$

Arguing by induction with respect to the index n, we deduce

$$\begin{aligned}
& \|(u^{(n+l+1)} - u^{(n+1)})(t)\|_{B^{s-1}_{p,r}} \\
\leq\ & C_T(2^{-n}\sum_{k=0}^{n}\frac{(2TC_T)^k}{k!} + C_T^{n+1}\int_0^t \frac{(t-\tau)^n}{n!}d\tau) \\
\leq\ & (C_T\sum_{k=0}^{n}\frac{(2TC_T)^k}{k!})2^{-n} + C_T\frac{(TC_T)^{n+1}}{(n+1)!},
\end{aligned}$$

which yields the desired result.

Finally, we can apply the interpolation method, which is similar to the proof in Lemma 3.1, to the critical case $s = 4 + \frac{1}{p}$. We here omit the details. Therefore, we complete the proof of Lemma 2.2. □

Based on the above preparations, we are in position to state the local existence result of the Cauchy problem (1.2).

Theorem 3.1. *Suppose that* $\leq p, r \leq \infty$, $s > \max\{2+\frac{1}{p}, \frac{5}{2}, 3-\frac{1}{p}\}$ *and* $u_0 \in B^s_{p,r}$. *Then there exists a time* $T > 0$ *such that the Cauchy problem (1.2) has a unique solution* $u \in E^s_{p,r}(T)$, *and the mapping* $u_0 \to u$ *is continuous from* $B^s_{p,r}$ *into*

$$C([0,T]; B^{s'}_{p,r}) \cap C^1([0,T]; B^{s'-1}_{p,r})$$

for all $s' < s$ *if* $r = \infty$ *and* $s' = s$ *otherwise.*

Proof. According to Lemma 3.1, $\{u^{(n)}\}_{n\in\mathbb{N}}$ is a Cauchy sequence in $C([0,T]; B^{s-1}_{p,r})$, so it converges to some function $u \in C([0,T]; B^{s-1}_{p,r})$. Thanks to Lemma 3.1 and Proposition 2.2 (iv) Fatou lemma, we have that $u \in L^\infty([0,T]; B^s_{p,r})$. Thus, by the interpolation method, for all $s' < s$, we find that $u \in C([0,T]; B^{s'}_{p,r})$.

Taking limit in Eq. (3.4), we conclude that u solves Eq. (1.2) in the sense of $u \in C([0,T]; B^{s'-1}_{p,r})$, for all $s' < s$. Since $u \in L^\infty([0,T]; B^s_{p,r})$ and the fact $B^s_{p,r}$ is an algebra as $s > 2 + \frac{1}{p}$, the right-hand side of the following equation

$$m_t - [\frac{k_1}{2}(u^2 - u_x^2) + \frac{k_2}{2}u]m_x = k_1 u_x m^2 + k_2 u_x m,$$

belongs to $L^\infty([0,T]; B^{s-2}_{p,r})$. In particular, for $r < \infty$, Lemma 2.2 enables us to get that $u \in C([0,T]; B^{s'}_{p,r})$ for all $s' < s$. Finally, taking advantage of Eq. (1.2) again, we obtain that $\partial_t u \in C([0,T]; B^{s-1}_{p,r})$ if $r < \infty$, and in $L^\infty([0,T]; B^{s-1}_{p,r})$ otherwise.

Moreover, the continuity with respect to the initial data in

$$C([0,T]; B^{s'}_{p,r}) \cap C^1([0,T]; B^{s'-1}_{p,r}) \quad (\forall s' < s)$$

can be obtained by Lemma 3.1 and a simple interpolation argument. While the case $s' = s$, a standard of use of a sequence of viscosity approximate solutions $\{u_\varepsilon\}_{\varepsilon>0}$ for Eq. (1.2) which converges uniformly in $C([0,T]; B^s_{p,r}) \cap C^1([0,T]; B^{s-1}_{p,r})$ gives the proof of the continuity of solutions in $E^s_{p,r}(T)$. This completes the proof of the theorem. □

Remark 3.1. *We know that nonhomogeneous Besov spaces contain Sobolev spaces. In fact, by Fourier-Plancherel formula, we find that the Besov space* $B^s_{2,2}(\mathbb{R})$ *coincides with the Sobolev space* $H^s(\mathbb{R})$. *Therefore, Theorem 3.1 implies that under the assumption* $u_0 \in H^s(\mathbb{R}), s > \frac{5}{2}$, *we can obtain the local well-posedness result to Eq. (1.2).*

Remark 3.2. *The existence time for Eq. (1.2) can be chosen independently of* s *in the following sense [30]. If*

$$u \in C([0,T]; H^s) \cap C^1([0,T]; H^{s-1})$$

is a solution to Eq. (1.2) with initial data $H^r, r > \frac{5}{2}, r \neq s$, *then*

$$u \in C([0,T]; H^r) \cap C^1([0,T]; H^{r-1})$$

with the same time $T > 0$. *In particular, if* $u_0 \in H^\infty$, *then* $u \in C([0,T]; H^\infty)$.

4 Blow-up scenario

In this section, by using the local well-posedness result of Theorem 3.1 and energy estimates, we present a precise blow-up scenario for strong solutions to the Cauchy problem (1.2).

Theorem 4.1. *Let $u_0 \in H^s(\mathbb{R}), s > \frac{5}{2}$ be given and assume that T is the maximal existence time of the solution $u(t,x)$ to Eq. (1.2) with the initial data u_0 guaranteed by Remark 3.1. Then the corresponding solution $u(t,x)$ blows up in finite time if and only if*

$$\liminf_{t\to T} \inf_{x\in\mathbb{R}} \{mu_x(t,x)\} = -\infty \qquad \textit{or} \qquad \liminf_{t\to T} \inf_{x\in\mathbb{R}} \{u_x(t,x)\} = -\infty.$$

Proof. From Remark 3.1-3.2 and a simple density argument, we only need to prove that Theorem 4.1 holds true for $s = 4$. Multiplying Eq. (1.2) by m, integrating over $\mathbb{R}$ and integration by parts, we get

$$\begin{aligned} & \frac{1}{2}\frac{d}{dt}\int_{\mathbb{R}} m^2 dx \qquad (4.1)\\ = \ & \frac{k_1}{2}\int_{\mathbb{R}}(u^2-u_x^2)mm_x dx + k_1\int_{\mathbb{R}} u_x m^3 dx + k_2\int_{\mathbb{R}} u_x m^2 dx + \frac{k_2}{2} umm_x dx \\ = \ & -\frac{k_1}{2}\int_{\mathbb{R}} u_x m^3 dx + k_1\int_{\mathbb{R}} u_x m^3 dx + k_2\int_{\mathbb{R}} u_x m^2 dx - \frac{k_2}{4}\int_{\mathbb{R}} u_x m^2 dx \\ = \ & \frac{k_1}{2}\int_{\mathbb{R}} u_x m^3 dx + \frac{3k_2}{4}\int_{\mathbb{R}} u_x m^2 dx. \end{aligned}$$

Differentiating Eq. (1.2) with respect to x, we deduce

$$\begin{aligned} m_{tx} &= 3k_1 u_x m m_x - k_1 m^3 + k_1 u m^2 + \frac{k_1}{2}(u^2 - u_x^2)m_{xx} \\ &\quad + \frac{3k_2}{2} m_x u_x + k_2 u m - k_2 m^2 + \frac{k_2}{2} u m_{xx}. \end{aligned}$$

Multiplying the above equation by m_x, and integrating with respect x over $\mathbb{R}$, we have

$$\begin{aligned} & \frac{1}{2}\frac{d}{dt}\int_{\mathbb{R}} m_x^2 dx \qquad (4.2)\\ = \ & 3k_1\int_{\mathbb{R}} u_x m m_x^2 dx - k_1\int_{\mathbb{R}} m^3 m_x dx + k_1\int_{\mathbb{R}} u m^2 m_x dx \\ & + \frac{k_1}{2}\int_{\mathbb{R}} (u^2 - u_x^2) m_x m_{xx} dx + \frac{3k_2}{2}\int_{\mathbb{R}} m_x^2 u_x dx \\ & + k_2\int_{\mathbb{R}} u m m_x dx - k_2\int_{\mathbb{R}} m^2 m_x dx + \frac{k_2}{2}\int_{\mathbb{R}} u m_x m_{xx} dx \\ = \ & 3k_1\int_{\mathbb{R}} u_x m m_x^2 dx - \frac{k_1}{3}\int_{\mathbb{R}} u_x m^3 dx - \frac{k_1}{4}\int_{\mathbb{R}} m_x^2 (u^2 - u_x^2)_x dx \\ & + \frac{3k_2}{2}\int_{\mathbb{R}} m_x^2 u_x dx - \frac{k_2}{2}\int_{\mathbb{R}} u_x m^2 dx - \frac{k_2}{4}\int_{\mathbb{R}} u - x m_x^2 dx \end{aligned}$$

$$= \frac{5k_1}{2}\int_{\mathbb{R}} u_x m m_x^2 dx - \frac{k_1}{3}\int_{\mathbb{R}} u_x m^3 dx + \frac{5k_2}{4}\int_{\mathbb{R}} m_x^2 u_x dx - \frac{k_2}{2}\int_{\mathbb{R}} u_x m^2 dx.$$

From (4.1)-(4.2), we get

$$\begin{aligned} &\frac{d}{dt}\int_{\mathbb{R}}(m^2+m_x^2)dx \\ = \; & 5k_1\int_{\mathbb{R}} u_x m m_x^2 dx + \frac{k_1}{3}\int_{\mathbb{R}} u_x m^3 dx + \frac{5k_2}{2}\int_{\mathbb{R}} m_x^2 u_x dx + \frac{k_2}{2}\int_{\mathbb{R}} u_x m^2 dx. \end{aligned}$$

Assume that $T < \infty$ and there exists $N_1, N_2 > 0$ such that $mu_x \geq -N_1, u_x \geq -N_2$ for all $(t,x) \in [0,T)\times\mathbb{R}$. Let us choose $N, k > 0$ such that $N := \max\{N_1, N_2\}$ and $k := \max\{-k_1, -k_2\}$. It then follows that

$$\frac{d}{dt}\int_{\mathbb{R}}(m^2+m_x^2)dx \leq 10kN\int_{\mathbb{R}}(m^2+m_x^2)dx.$$

Applying Gronwall's lemma to the above inequality implies for $t \in [0,T)$,

$$\|m\|_{H^1}^2 \leq e^{10NkT}\|m_0\|_{H^1}^2. \tag{4.3}$$

Differentiating Eq. (1.2) with respect to x twice, we deduce

$$\begin{aligned} m_{txx} = \; & -6k_1 m^2 m_x + 5k_1 u m m_x + 4k_1 u_x m m_{xx} + 3k_1 u_x m_x^2 \\ & + k_1 u_x m^2 + \frac{k_1}{2}(u^2-u_x^2)m_{xxx} + 2k_2 u_x m_{xx} - \frac{7k_2}{2} m m_x \\ & + \frac{5k_2}{2} u m_x + k_2 u_x m + \frac{k_2}{2} u m_{xxx}. \end{aligned}$$

Multiplying the above equation by m_{xx}, integrating with respect to x over $\mathbb{R}$, we have

$$\begin{aligned} & \frac{1}{2}\frac{d}{dt}\int_{\mathbb{R}} m_{xx}^2 dx \qquad (4.4) \\ = \; & -6k_1\int_{\mathbb{R}} m^2 m_x m_{xx} dx + 5k_1\int_{\mathbb{R}} u m m_x m_{xx} dx + 4k_1\int_{\mathbb{R}} u_x m m_{xx}^2 dx \\ & + 3k_1\int_{\mathbb{R}} u_x m_x^2 m_{xx} dx + k_1\int_{\mathbb{R}} u_x m^2 m_{xx} dx + \frac{k_1}{2}\int_{\mathbb{R}}(u^2-u_x^2)m_{xx}m_{xxx} dx \\ & + 2k_2\int_{\mathbb{R}} u_x m_{xx}^2 dx - \frac{7k_2}{2}\int_{\mathbb{R}} m m_x m_{xx} dx + \frac{5k_2}{2}\int_{\mathbb{R}} u m_x m_{xx} dx \\ & + k_2\int_{\mathbb{R}} u_x m m_{xx} dx + \frac{k_2}{2}\int_{\mathbb{R}} u m_{xx} m_{xxx} dx \\ = \; & -6k_1\int_{\mathbb{R}} m^2 m_x m_{xx} dx + 5k_1\int_{\mathbb{R}} u m m_x m_{xx} dx + 4k_1\int_{\mathbb{R}} u_x m m_{xx}^2 dx \\ & + k_1\int_{\mathbb{R}} u_x m m_x^2 dx + 2k_1\int_{\mathbb{R}} u m m_x m_{xx} dx - k_1\int_{\mathbb{R}} m^2 m_x m_{xx} dx \\ & - 2k_1\int_{\mathbb{R}} u_x m m_x^2 dx + \frac{k_1}{3}\int_{\mathbb{R}} u_x m^3 dx - \frac{k_1}{2}\int_{\mathbb{R}} u_x m m_{xx}^2 dx \\ & + 2k_2\int_{\mathbb{R}} u_x m_{xx}^2 dx - \frac{7k_2}{2}\int_{\mathbb{R}} m m_x m_{xx} dx - \frac{5k_2}{4}\int_{\mathbb{R}} u_x m_x^2 dx \end{aligned}$$

$$
\begin{aligned}
&-k_2\int_{\mathbb{R}}u_xm_x^2dx+\frac{k_2}{2}\int_{\mathbb{R}}u_xm^2dx-\frac{k_2}{4}\int_{\mathbb{R}}u_xm_{xx}^2dx\\
=\;&-7k_1\int_{\mathbb{R}}m^2m_xm_{xx}dx+7k_1\int_{\mathbb{R}}umm_xm_{xx}dx+\frac{7k_1}{2}\int_{\mathbb{R}}u_xmm_{xx}^2dx\\
&-k_1\int_{\mathbb{R}}u_xmm_x^2dx+\frac{k_1}{3}\int_{\mathbb{R}}u_xm^3dx+\frac{7k_2}{4}\int_{\mathbb{R}}u_xm_{xx}^2dx\\
&-\frac{7k_2}{2}\int_{\mathbb{R}}mm_xm_{xx}dx-\frac{9k_2}{4}\int_{\mathbb{R}}u_xm^2dx+\frac{k_2}{2}\int_{\mathbb{R}}u_xm^2dx.
\end{aligned}
$$

Combining (4.1)-(4.2) and (4.4), we obtain

$$
\begin{aligned}
&\frac{d}{dt}\int_{\mathbb{R}}(m^2+m_x^2+m_{xx}^2)dx\\
=\;&-14k_1\int_{\mathbb{R}}m^2m_xm_{xx}dx+14k_1\int_{\mathbb{R}}umm_xm_{xx}dx+7k_1\int_{\mathbb{R}}u_xmm_{xx}^2dx\\
&+3k_1\int_{\mathbb{R}}u_xmm_x^2dx+k_1\int_{\mathbb{R}}u_xm^3dx+\frac{7k_2}{2}\int_{\mathbb{R}}u_xm_{xx}^2dx\\
&-7k_2\int_{\mathbb{R}}mm_xm_{xx}dx-2\int_{\mathbb{R}}u_xm^2dx+\frac{3k_2}{2}\int_{\mathbb{R}}u_xm^2dx.
\end{aligned}
$$

If mu_x and u_x are bounded from below on $[0,T)\times\mathbb{R}$, *i.e.*, there exists $N_1,N_2>0$ such that $mu_x\geq -N_1, u_x\geq -N_2$ for all $(t,x)\in[0,T)\times\mathbb{R}$. Similarly, we can choose $N,k>0$ such that $N:=\max\{N_1,N_2\}$ and $k:=\max\{-k_1,-k_2\}$. Then, by (4.3) and the above equality, we get

$$
\begin{aligned}
&\frac{d}{dt}\int_{\mathbb{R}}(m^2+m_x^2+m_{xx}^2)dx\\
\leq\;&\frac{21}{2}kN\int_{\mathbb{R}}(m^2+m_x^2+m_{xx}^2)dx+14k(\|m\|_{L^\infty}^2+\|um\|_{L^\infty}\\
&+\|m\|_{L^\infty})\int_{\mathbb{R}}|m_xm_{xx}|dx\\
\leq\;&\frac{21}{2}kN\int_{\mathbb{R}}(m^2+m_x^2+m_{xx}^2)dx+14k(\|m\|_{H^1}^2+\|m\|_{H^1})\int_{\mathbb{R}}(m_x^2+m_{xx}^2)dx\\
\leq\;&7k[\frac{3N}{2}+2e^{5NkT}\|m_0\|_{H^1}(e^{5NkT}\|m_0\|_{H^1}+1)]\int_{\mathbb{R}}(m^2+m_x^2+m_{xx}^2)dx.
\end{aligned}
$$

Hence, applying Gronwall's inequality implies that for all $t\in[0,T)$

$$
\begin{aligned}
\|u\|_{H^4}^2\;&\leq\;\|m\|_{H^2}^2=\int_{\mathbb{R}}(m^2+m_x^2+m_{xx}^2)dx\\
&\leq\;\exp\{7k[\frac{3N}{2}+2e^{5NkT}\|m_0\|_{H^1}(e^{5NkT}\|m_0\|_{H^1}+1)]\}\|m_0\|_{H^2}^2\\
&\leq\;C\exp\{7k[\frac{3N}{2}+2e^{5NkT}\|m_0\|_{H^1}(e^{5NkT}\|m_0\|_{H^1}+1)]\}\|u_0\|_{H^4}^2.
\end{aligned}
$$

The above inequality and Sobolev's embedding theorem ensure that $u(t,x)$ does not blow up in finite time.

On the other hand, by Sobolev's imbedding theorem, we find that if $\lim\inf_{t\to T}\inf_{x\in\mathbb{R}}\{mu_x(t,x)\} = -\infty$ or $\lim\inf_{t\to T}\inf_{x\in\mathbb{R}}\{u_x(t,x)\} = -\infty$, then the solution will blow up in finite time. This completes the proof of the theorem. □

5 Peakons and weak kinks and their interactions

Let us consider the following equation [26]

$$m_t = bu_x + \frac{1}{2}k_1\left[m(u^2-u_x^2)\right]_x + \frac{1}{2}k_2(2mu_x+m_xu), \quad m = u-u_{xx}, \tag{5.1}$$

where b, k_1 and k_2 are three arbitrary constants. Apparently, when $b=0$, equation (5.1) recovers Eq. (1.2).

To analyze peakons for the nonlinear cubic equation (5.1), let us first investigate the weak form of the equation. Applying the operator $(1-\partial_x^2)^{-1}$ to equation (5.1), we obtain

$$\begin{aligned} u_t &= \frac{1}{6}k_1(3u^2u_x-u_x^3)+\frac{1}{2}k_2uu_x+\frac{1}{6}k_1(1-\partial_x^2)^{-1}u_x^3 \\ &\quad +\frac{1}{2}\partial_x(1-\partial_x^2)^{-1}\left(2bu+k_1(uu_x^2+\frac{2}{3}u^3)+k_2(u^2+\frac{1}{2}u_x^2)\right). \end{aligned} \tag{5.2}$$

Taking the convolution with the Green's function $G(x)=\frac{1}{2}\exp(-|x|)$ for the Helmholtz operator $(1-\partial_x^2)$, equation (5.2) can be rewritten as

$$\begin{aligned} u_t &= \frac{1}{6}k_1(3u^2u_x-u_x^3)+\frac{1}{2}k_2uu_x+\frac{1}{6}k_1G(x)*u_x^3 \\ &\quad +\frac{1}{2}\partial_x\left(G(x)*[2bu+k_1(uu_x^2+\frac{2}{3}u^3)+k_2(u^2+\frac{1}{2}u_x^2)]\right). \end{aligned} \tag{5.3}$$

When $b=0$, the following function

$$u = Ce^{-|x-ct|}, \quad C = \frac{-3\left(\sqrt{3}k_2 \pm \sqrt{3k_2^2-16k_1c}\right)}{4\sqrt{3}k_1} \tag{5.4}$$

satisfies equation (5.3), and therefore this is a single peakon solution of equation (1.2). Apparently, if $3k_2^2-16k_1c\geq 0$, then (5.4) gives a real peakon solution, and if $3k_2^2-16k_1c<0$, then we obtain a complex peakon. So, here we propose the complex peakon solutions for integrable equation (1.2), which is another difference from the CH equation.

Let us now consider the two-peakon solution to equation (1.2) in the following form

$$u = p_1(t)e^{-|x-q_1(t)|}+p_2(t)e^{-|x-q_2(t)|}. \tag{5.5}$$

Substituting (5.5) into Eq. (1.2), we may obtain the following peakon dynamical system

$$\begin{cases} p_{1,t} = -\frac{1}{2}k_2 p_1 p_2 sgn(q_1 - q_2) e^{-|q_1-q_2|}, \\ p_{2,t} = -\frac{1}{2}k_2 p_1 p_2 sgn(q_2 - q_1) e^{-|q_2-q_1|}, \\ q_{1,t} = -k_1 p_1 p_2 e^{-|q_1-q_2|} - \frac{1}{3}k_1 p_1^2 - \frac{1}{2}k_2 \left(p_1 + p_2 e^{-|q_1-q_2|}\right), \\ q_{2,t} = -k_1 p_1 p_2 e^{-|q_1-q_2|} - \frac{1}{3}k_1 p_2^2 - \frac{1}{2}k_2 \left(p_2 + p_1 e^{-|q_1-q_2|}\right). \end{cases} \tag{5.6}$$

For $k_1 = 0$, $k_2 = -2$, (1.2) recovers the CH equation and (5.6) is reduced to the two-peakon dynamics of CH equation, which Camassa and Holm studied in [3].

For $k_2 = 0$ and $b = 0$, (1.2) is exactly the cubic nonlinear equation (1.1) and the peakon dynamical system (5.6) is simplified to

$$\begin{cases} p_{1,t} = 0, \quad p_{2,t} = 0, \\ q_{1,t} = -k_1 p_1 p_2 e^{-|q_1-q_2|} - \frac{1}{3}k_1 p_1^2, \\ q_{2,t} = -k_1 p_1 p_2 e^{-|q_1-q_2|} - \frac{1}{3}k_1 p_2^2. \end{cases} \tag{5.7}$$

For the case of $c_1^2 = c_2^2$, the solutions of $q_1(t)$ and $q_2(t)$ in (5.7) are given by

$$\begin{cases} q_1(t) = -k_1(c_1 c_2 e^{-|A_1-A_2|} + \frac{1}{3}c_1^2)t + A_1, \\ q_2(t) = -k_1(c_1 c_2 e^{-|A_1-A_2|} + \frac{1}{3}c_1^2)t + A_2, \end{cases} \tag{5.8}$$

where A_1, A_2 are arbitrary integral constants. If $A_1 = A_2$, the two-peakon solution is reduced to zero (when $c_2 = -c_1$) or a single-peakon (when $c_2 = c_1$). If $A_1 \neq A_2$, the two-peakon can not collide since $q_1(t) \neq q_2(t)$ for every $t \in (-\infty, +\infty)$. Especially, in the peakon-antipeakon case ($c_1 = -c_2$), the collision can't occur for $A_1 \neq A_2$. This is very different from the case of the CH equation [3]. For the case of $c_1^2 \neq c_2^2$, we obtain the following solution:

$$\begin{cases} p_1(t) = c_1, \quad p_2(t) = c_2, \\ q_1(t) = sgn(t)\frac{3k_1c_1c_2}{|k_1(c_1^2-c_2^2)|}\left(e^{-|\frac{1}{3}k_1(c_1^2-c_2^2)t|} - 1\right) - \frac{1}{3}k_1 c_1^2 t, \\ q_2(t) = sgn(t)\frac{3k_1c_1c_2}{|k_1(c_1^2-c_2^2)|}\left(e^{-|\frac{1}{3}k_1(c_1^2-c_2^2)t|} - 1\right) - \frac{1}{3}k_1 c_2^2 t. \end{cases} \tag{5.9}$$

So, our results show that the collision of two-peakon of equation (1.1) is very different from the case of CH equation. In the peakon-antipeakon case, the collision can occur for the CH equation, but could not happen for equation (1.1). The collision of two-peakon of equation (1.1) occurs in the "chase" case as shown above.

For the case $k_1 \neq 0$ and $k_2 \neq 0$, let us choose $k_1 = k_2 = -2$. Then after a longer computation for Eq. (5.6), we arrive at

$$\begin{aligned} p_1(t) &= \coth(t), \quad q_1(t) = \frac{8}{3(e^{2t}-1)} + \ln(e^{2t}+1) - \frac{1}{3}t - \ln 2, \\ p_2(t) &= -\coth(t), \quad q_2(t) = \frac{8}{3(e^{2t}-1)} - \ln(e^{2t}+1) + \frac{5}{3}t + \ln 2, \end{aligned} \tag{5.10}$$



Figure 1: The peakon-antipeakon solution (5.11). Pink: peakon (and antipeakon) with the peak (and trough) position q_2; Green: antipeakon (and peakon) with the trough (and peak) position q_1.



Figure 2: The kink-peakon interacted solution. Black line: $t = 2$; Blue line: $t = 1$; Green line: $t = 0$.

which leads to the following peakon-antipeakon solution

$$u(x,t) = \coth(t)\left(e^{-|x-q_1(t)|} - e^{-|x-q_2(t)|}\right). \tag{5.11}$$

See Figure 1 for the profile of the peakon-antipeakon solution.

We have already shown that equation (5.1) admits peakon solutions in the case of $b = 0$. It is natural to ask what kind of solution one may obtain for the case of $b \neq 0$. Here we will reveal that equation (5.1) with $k_2 = 0$ and $b \neq 0$ possesses the weak kink and kink-peakon interacted solutions. Actually, equation (5.1) has the following weak kink solution

$$u = Csgn(x-ct)\left(e^{-|x-ct|} - 1\right), \quad C = \pm\sqrt{\frac{-b}{k_1}} \tag{5.12}$$

where the conditions $k_2 = 0, k_1 \neq 0, b \neq 0, c = -\frac{1}{2}b$ are required for equation (5.1). u is called a **weak kink solution** since u, u_x, u_t are continuous and u_{xx} has a jump at the peak point. In particular, we take $k_2 = 0$, $k_1 = -b = 2$, then equation (5.1) is cast to

$$m_t + 2u_x - \left[m(u^2 - u_x^2)\right]_x = 0, \quad m = u - u_{xx}, \tag{5.13}$$

and the corresponding weak kink solution is

$$u = sgn(x-t)\left(e^{-|x-t|} - 1\right). \tag{5.14}$$

Let us assume the following ansatz of kink-peakon interacted solutions

$$u = p_1(t)sgn(x-q_1(t))\left(e^{-|x-q_1(t)|} - 1\right) + p_2(t)e^{-|x-q_2(t)|}, \tag{5.15}$$

which actually describes **a new phenomena of kink-peakon interacted dynamics** in soliton theory. Substituting (5.15) into (5.3) with $k_2 = 0$, we have

$$\begin{cases} p_1 = \pm\sqrt{\frac{-b}{k_1}}, \\ p_{2,t} = k_1p_1^2p_2sgn(q_2 - q_1)e^{-|q_1-q_2|}, \\ q_{1,t} = -\frac{1}{2}b - k_1p_1p_2sgn(q_2 - q_1)e^{-|q_1-q_2|}, \\ q_{2,t} = -\frac{1}{3}k_1p_2^2 - \frac{1}{2}k_1p_1^2 + k_1\left(p_1^2 - p_1p_2sgn(q_2 - q_1)\right)e^{-|q_1-q_2|} + k_1sgn(q_2 - q_1)p_1p_2. \end{cases} \tag{5.16}$$

Choosing $k = -b = 2, p_1 = 1$ and $q_1 < q_2$, we may obtain the following exact solution:

$$\begin{cases} q_1 = t - p_2, \\ q_2 = t - p_2 - \ln(\frac{1}{9}p_2^2 - \frac{1}{2}p_2 + 1 + \frac{A_2}{2p_2}), \\ p_{2,t} = \frac{2}{9}p_2^3 - p_2^2 + 2p_2, \end{cases} \tag{5.17}$$

and p_2 is able to be solved in the following implicit form:

$$\ln|p_2| - \frac{1}{2}\ln(p_2^2 - \frac{9}{2}p_2 + 9) + \frac{3\sqrt{7}}{7}\arctan(\frac{4p_2 - 9}{3\sqrt{7}}) = 2t. \tag{5.18}$$

See Figure 2 for the profile of the kink-peakon interactional solution (5.15) with $k = -b = 2, p_1 = 1$.

In general, we may study the N-peakon system for more detailed investigations, and also we may make the following ansatz of the solution for equation (5.1) with $k_2 = 0$ and $b \neq 0$:

$$u = p_0(t)sgn(x - q_0(t))\left(e^{-|x-q_0(t)|} - 1\right) + \sum_{j=1}^{N} p_j(t)e^{-|x-q_j(t)|}, \tag{5.19}$$

which can be viewed as the interaction of single-kink and N-peakon solution. How we handle such interactions will be our future work.

Acknowledgments Yin was partially supported by NNSFC (No. 11271382 and No. 10971235), RFDP (No. 20120171110014), and the key project of Sun Yat-sen University (No. c1185). Qiao was partially supported by the Texas Norman Hackerman Advanced Research Program (Grant No. 003599-0001-2009) and supervised the U.S. Department of Education UTPA-GAANN program (P200A120256).

References

[1] H. Bahouri, Y. Y. Chemin and R. Danchin, Fourier Analysis and Nonlinear Pratial Differential Equations, Grundlehren der mathematischen Wissenschaften, Springer, 2011, pages 51-167.

[2] P. M. Bies, P. Górka and E. Reyes, The dual modified Korteweg-de Vries-Fokas-Qiao equation: Geometry and local analysis, *J. Math. Phys.,* **53** (2012), 073710.

[3] R. Camassa and D. Holm, An integrable shallow water equation with peaked solitons, *Phys. Rev. Lett.,* **71** (1993), 1661-1664.

[4] R. Camassa, D. Holm and J. Hyman, A new integrable shallow water equation, *Adv. Appl. Mech.,* **31** (1994), 1-33.

[5] A. Constantin, Global existence of solutions and breaking waves for a shallow water equation: a geometric approach, *Ann. Inst. Fourier(Grenoble),* **50** (2000), 321-362.

[6] A. Constantin and J. Escher, Wave breaking for nonlinear nonlocal shallow water equations, *Acta Math.,* **181** (1998), 229-243.

[7] A. Constantin and J. Escher, Well-posedness, global existence and blowup phenomena for a periodic quasi-linear hyperbolic equation, *Comm. Pure Appl. Math.,* **51** (1998), 475-504.

[8] R. Danchin, A few remarks on the Camassa-Holm equation, *Diff. Int. Eq.,* **14** (2001), 953-988.

[9] J. A. Desanto, Mathematical and numerical aspects of wave propagation: proceedings in applied mathematics, (Proceedings in Applied Mathematics Series Siam Proceedings Series), 1998.

[10] A. Fokas, The Korteweg-de Vries equation and beyond, *Acta Appl. Math.,* **39** (1995), 295-305.

[11] A. Fokas and B. Fuchssteiner, Symplectic structures, their Bäcklund transformation and hereditary symmetries, *Physica D,* **4** (1981), 47-66.

[12] B. Fuchssteiner, Some tricks from the symmetry-toolbox for nonlinear equations: Generalizations of the Camassa-Holm equation, *Physica D,* **95** (1996), 229-243.

[13] G. Gui, Y. Liu, P. J. Olver and C. Qu, Wave-breaking and peakons for a modified Camassa-Holm equation, *Comm. Math. Phys.,* (2012), DOI: 10.1007/s00220-012-1566-0.

[14] A. Himonas and C. Holliman, The Cauchy problem for the Novikov equation, *Nonlinearity,* **25** (2012), 449-479.

[15] A. N. Hone, H. Lundmark and J. Szmigielski, Explicit multipeakon solutions of Novikovs cubically nonlinear integrable Camassa-Holm type equation, *Dyn. Partial Diff. Eqns.,* **6** (2009), 253-89.

[16] A. N. Hone and J. Wang, Integrable peakon equations with cubic nonlinearity, *J. Phys. A: Math. Theor.,* **41** (2008), 372002.

[17] J. Lenells, Traveling wave solutions of the Camassa-Holm equation, *J. Differential Equations,* **217** (2005), 393-430.

[18] J. B. Li and Y. Zhang, Exact M/W-shape solitary wave solutions determined by a singular traveling wave equation, *Nonlinear Anal. Real World Appl.*, **10** (2009), 1797-1802.

[19] Y. Li and P. J. Olver, Well-posedness and blow-up solutions for an integrable nonlinear dispersive model wave equation, *J. Differential Equations,* **162** (2000), 27-63.

[20] X. Liu and Z. Yin, Low regularity solutions, blow up, and global existence for a generalization of Camassa-Holm-type equation, submitted.

[21] V. Novikov, Generalizations of the Camassa-Holm equation, *J. Phys. A: Math. Theor.,* **42** (2009), 342002.

[22] P. J. Olver and P. Rosenau, Tri-Hamiltonian duality between solitons and solitary-wave solutions having compact support, *Phys. Rev. E,* **53** (1996), 1900-1906.

[23] A. R. Osborne, 33-modeling the 2+1 Gardner equation, *Int. Geophys,* **97** (2010), 857-866.

[24] Z. Qiao, A new integrable equation with cuspon and W/M-shape-peaks solitons, *J. Math. Phys.,* **47** (2006), 112701; New integrable hierarchy, parametric solutions, cuspons, one-peak solitons, and M/W-shape solutions, *J. Math. Phys.,* **48** (2007), 082701.

[25] Z. Qiao and X. Li, An integrable equation with nonsmooth solitons, *Theor. Math. Phys.,* **267** (2011), 584-589.

[26] Z. Qiao, B. Xia and J. B. Li, A generalized Camassa-Holm equation and its peakon solutions, http://xxx.lanl.gov/abs/1205.2028.

[27] F. Tiğlay, The periodic Cauchy problem for Novikovs equation, *Int. Math. Res. Not.,* **20** (2011), 4633-48.

[28] X. Wu and Z. Yin, Global weak solutions for the Novikov equation, *J. Phys. A: Math. Theor.,* **44** (2011), 055202.

[29] X. Wu and Z. Yin, A note on the Cauchy problem of the Novikov equation, *Appl. Anal.,* (2012), DOI:10.1080/00036811.2011.649735.

[30] Z. Yin, On the Cauchy problem for an integrable equation with peakon solutions, *Illinois. J. Math.,* **47** (2003), 649-666.